

(Due Wednesday 09/19/2018 right before class)

&

(Your homework shall be stapled if it contains multiple pages.)

**FALL/2018/MA526: HOMEWORK 4**

Instructor: Guangqu Zheng<sup>1</sup>; Grader: Chessa Mccalla<sup>2</sup>

**Total points: 20.**

**Q1** Students shall show enough details for obtaining their answers.

(1) Find the value of  $C$  such that  $f$  defined below is a probability density function: (2 points)

$$f(x) = \begin{cases} C \cdot x^{-3} & \text{if } x \geq 2 \\ 0 & \text{if } x < 2. \end{cases}$$

(2) Find the value of  $C$  such that  $g$  defined below is a probability density function: (3 points)

$$g(x) = \begin{cases} C \cdot x^{-3} & \text{if } x \geq 2 \\ C \cdot x^{-2} & \text{if } 1 \leq x < 2. \\ 0 & \text{if } x < 1. \end{cases}$$

(3) Find the values of  $A$  and  $B$  such that  $h$  defined below is a probability density function: (3 points)

$$h(x) = \begin{cases} Ae^{-x} & \text{if } x \geq 1 \\ Bx & \text{if } -1 \leq x \leq 1. \\ \frac{1}{2}e^x & \text{if } x \leq -1. \end{cases}$$

(4) Solve **exercise 3.12 page 92** in the *textbook* (see the following screenshot) (4 points)

**3.12** An investment firm offers its customers municipal bonds that mature after varying numbers of years. Given that the cumulative distribution function of  $T$ , the number of years to maturity for a randomly selected bond, is

$$F(t) = \begin{cases} 0, & t < 1, \\ \frac{1}{4}, & 1 \leq t < 3, \\ \frac{1}{2}, & 3 \leq t < 5, \\ \frac{3}{4}, & 5 \leq t < 7, \\ 1, & t \geq 7, \end{cases}$$

find

- (a)  $P(T = 5)$ ;
- (b)  $P(T > 3)$ ;
- (c)  $P(1.4 < T < 6)$ ;
- (d)  $P(T \leq 5 \mid T \geq 2)$ .

---

<sup>1</sup>gzheng90@ku.edu; Office hours: MWF 3:00-3:50; Office = 641 Snow Hall

<sup>2</sup>chessa\_m@ku.edu

**Q2** (4 points) Let  $W$  be a random variable giving the number of heads minus the number of tails in three tosses of a coin.

- 1) List the elements of the sample space  $\Omega$  for the three tosses of the coin. (1 point)
- 2) Assume that the probability of getting a head is equal to  $1/2$ . Find out the probability mass function  $f$  associated to  $W$ . (2 points)
- 3) Find out the cumulative distributional function associated to the pmf  $f$  given in part-2. (1 point)

**Q3** (4 points) Let  $A, B, C$  be three events on some sample space  $\Omega$  such that  $A$  and  $B$  are independent while  $(A \cup B) \cap C = \emptyset$ . Assume that  $\mathbb{P}(C) = 0.1$ ,  $\mathbb{P}(A) = 0.5$  and  $\mathbb{P}(B) = 0.4$ .

We define now

$$X = \mathbb{I}_A + \mathbb{I}_B + \mathbb{I}_C.$$

- (a) Prove  $A$  and  $C$  are not independent. (1 point)      (b) Find the CDF of  $X$ . (3 points)

Recall  $\mathbb{I}_A$  is the indicator function over  $A$ , defined as follows:

$$\mathbb{I}_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$